



MALARIA CAREMAP

Diagnostic algorithm of malaria¹⁻⁴



RDT: Rapid diagnostic test

Management algorithm of malaria^{1,2,4,5}



*Doxycycline is contraindicated in pregnant and lactating women and children less than 8 years; ACT: Artemisinin combination therapy; CQ: Chloroquine; IM: Intramuscular; IV: Intravenous; PQ: Primaquine

References: 1. World Health Organization. Guidelines for the treatment of malaria 2015 (3rd edition). Available from: https://www.who.int/docs/default-source/documents/publications/gmp/guidelines-for-the-treatment-of-malaria-eng.pdf?sfvrsn=a0138b77_2. Last accessed on 1st June 2019. 2. National Institute of Malaria Research. Guidelines for diagnosis and treatment of malaria in India 2014 (3rd edition). Available from: <https://nvbdcp.gov.in/WriteReadData/1892s/20627628441542176662.pdf>. Last accessed on 1st June 2019. 3. Mathison BA, et al. J Clin Microbiol. 2017;55(7):2009-17. 4. Centers for Disease Control and Prevention. Malaria treatment and diagnosis. Available from: https://www.cdc.gov/malaria/diagnosis_treatment/treatment.html. Last accessed on 1st June 2019. 5. Tan KR, et al. Am J Trop Med Hyg. 2011;84(4):517-31.

DOXTTM Injection
Doxycycline hyclate 100 mg